



Aesthetic Clinics & Hair Salon

The only one hair mesotherapy solution
approved by SCI journal & 10 Univ.Hospitals

User Manual

HR CYTOKINES

SGF 57



Recommended program in clinics (HR Cytokines Therapy)

Forte (High concentration)
2 vials
 7 cytokines about
 20.5ppm / 5 cc diluted per vial



Basic (Standard concentration)
8 vials
 7 cytokines about
 5.5ppm / 5cc diluted per vial

1 Box contains 12 treatments for 2 person : Total 24 treatments

3Months(Total 12 treatments)

Week 1	Week 2	Week 3	Week 4	Week 5	Week 6	Week 7	Week 8	Week 9	Week 10	Week 11	Week 12
Forte	Forte	Forte	Forte	Basic	Basic	Basic	Basic	Basic		Basic	

- Intensive treatment with 1 Forte vial for 4 weeks
- 1.25 cc per session


- From Week 5 through Week 8:
Treatment with basic vial
- 1 time per Week
- 2.5 cc per session

- From 9 week through 12week: Depending on the progress of the treatment, treatment intervals can be changed to 2 or 4 Weeks interval
- Treatment with basic vial
- 1times every 2 Weeks or 4 Weeks
- 2.5 cc per session

Recommended direction of use

Step 1

Prepare.....



Recommended direction of use

Open the
Normal
saline

Step 2



Recommended direction of use

Extract 5ml of
Normal saline
with syringe
needle



Recommended direction of use

Open the
SGF-57 vial
cap

Step 4



Recommended direction of use

Insert syringe
needle into
SGF-57 vial



Note : Normal saline will automatically go into the SGF-57 vial since vial is in vacuum state

Recommended direction of use

Shake the
SGF-57 vial



Recommended direction of use

When using,
Forte : Extract 1.25cc
Basic : Extract 2.5cc of
SGF-57 into the syringe
needle

Note : After extraction, store the SGF-57 mixed with saline in the refrigerator. From our trial test, it was found that the saline mixed SGF57's cytokines activity is above 80% when used within 3 months



Recommended direction of use

Insert the
extracted mixture
of SGF-57 into the
vial tube

Note : If the scalp area to be treated is
wide....just add more saline.
For instance, Forte 1.25cc+more saline.
Basic 2.5cc+more saline

Step 8



SGF-57 Recommended direction of use

Apply SGF-57 on the scalp by using MTS 0.2~1.0mm needle depth





SGF-57 Recommended direction of use

Use electroporation device to infiltrate the SGF-57 further and massage the scalp (if available)

Step 10



Step by Step process HR cytokines treatment for clinics(Hair)



Post procedure treatment

- After treatment prescribe anti-hair loss shampoo and Trico Clin tonic
- Advice patients on home-care practices on slowing hair loss



HR CYTOKINES

SGF-57

The only one hair
mesotherapy solution approved by
SCI journal & 10 Univ.Hospitals


AESMED



Thank You
→